



Waterfowl Breeder Diet

Mazuri® Waterfowl Breeder Diet contains biologically available nutrients to enhance breeding performance, egg numbers, fecundity, and chick viability for ornamental waterfowl.

Features and Benefits

- **Exceeds published niacin requirements for waterfowl** – Supports healthy growth and development.
- **Nutritionally balanced** – Promotes successful breeding programs and requires no mixing or supplementation.
- **Floating, extruded diet** – Allows birds to be fed on the water and prevents accumulation on the bottom of the feeding area, limiting water contamination and waste and creating a more natural feeding environment.
- **Sustainable fishmeal source used.**

Product Form & Packaging

Catalog #0001433 | Extruded Particle | 3/8" x 3/16" | 50 lb. (22.67 kg) paper w/poly bag

Guaranteed Analysis

Crude protein not less than.....	17.00%	Calcium not more than	3.50%
Crude fat not less than	2.50%	Phosphorus not less than.....	0.70%
Crude fiber not more than	6.00%	Salt not less than.....	0.05%
Ash not more than	11.00%	Salt not more than	0.55%
Calcium not less than	2.50%	Sodium not more than	0.55%

Ingredients

Ground Corn, Dehulled Soybean Meal, Wheat Middlings, Calcium Carbonate, Fish Meal, Dehydrated Alfalfa Meal, Soybean Oil, Dicalcium Phosphate, Salt, Brewers Dried Yeast, Choline Chloride, Cholecalciferol (Vitamin D3), Menadione Sodium Bisulfite Complex (Vitamin K), Vitamin A Acetate, Biotin, Manganous Oxide, d-Alpha Tocopheryl Acetate (Vitamin E), Zinc Oxide, Folic Acid, Calcium Pantothenate, Nicotinic Acid, Pyridoxine Hydrochloride, Copper Sulfate, Vitamin B12 Supplement, Riboflavin Supplement, Preserved with Mixed Tocopherols, Magnesium Oxide, Rosemary Extract, Citric Acid (a Preservative), DL-Methionine, Thiamine Mononitrate, L-Lysine, Calcium Iodate, Sodium Selenite.

Feeding Directions

- This diet can be fed to birds as a major part of the daily food intake during the breeding season (spring months).
- Feed free choice to mature birds 2 to 4 weeks before breeding season.
- Adult birds should eat in the range of 25 to 30 gram per kilogram of body weight each day, depending on species size.
 - Supplementation is not necessary, but if desired, other food items and supplements should make up no more than 20% of the total dietary intake.
- To encourage natural feeding behavior, scatter food directly on the water, where it will float.
 - To prevent the diet from floating away, offer feed in a shallow trough of water.
- For added convenience, this diet may also be offered dry.
- Always provide plenty of fresh, clean water.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

Storage Conditions

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.