



Waterfowl Starter

Mazuri® Waterfowl Starter provides appropriate nutrition for the rapid growth requirements of newly hatched chicks and adolescent birds. This diet can be fed by floating on the water or in a trough.

Features and Benefits

- **High protein, nutrient-dense diet** – Supports health and growth.
- **Exceeds published niacin requirements for waterfowl** – Supports healthy growth and development at 150 ppm.
- **Floating, extruded diet** – Allows birds to be fed on the water and prevents accumulation on the bottom of the feeding area, limiting water contamination and waste and creating a more natural feeding environment.
- **No supplementation required** – Requires no mixing or supplementation and eliminates sorting.
- **Sustainable fishmeal source used.**

Product Form & Packaging

Extruded Particle | 3/32" x 3/32"

- **Catalog #0057007** | 5 lb. (2.26 kg) net weight resealable poly bag
- **Catalog #0001434** | 25 lb. (11.33 kg) net weight woven poly bag
- **Catalog #0057008** | Case: 4 x 5 lb. net weight resealable poly bags

Guaranteed Analysis

Crude protein not less than	20.00%	Calcium not more than	1.45%
Crude fat not less than	3.00%	Phosphorus not less than	0.85%
Crude fiber not more than	6.50%	Salt not less than	0.05%
Ash not more than	7.00%	Salt not more than	0.55%
Calcium not less than	0.95%	Niacin not less than	70 mg/lb

Ingredients

Ground Corn, Wheat Middlings, Dehulled Soybean Meal, Ground Wheat, Fish Meal, Soybean Oil, Calcium Carbonate, Dicalcium Phosphate, Brewers Dried Yeast, Salt, DL-Methionine, Cholecalciferol (Vitamin D3), d-Alpha Tocopheryl Acetate (Vitamin E), Pyridoxine Hydrochloride, Menadione Sodium Bisulfite Complex (Vitamin K), Preserved with Mixed Tocopherols, Nicotinic Acid, Vitamin A Acetate, Choline Chloride, Manganous Oxide, Citric Acid (a Preservative), Zinc Oxide, Calcium Pantothenate, Folic Acid, Tagetes (Aztec Marigold) Extract (Color), Copper Sulfate, Thiamine Mononitrate, Riboflavin-5-Phosphate, Vitamin B12 Supplement, L-Lysine, Calcium Iodate, Rosemary Extract, Sodium Selenite, Biotin

Feeding Directions

- Feed free choice to newly hatched and adolescent birds during their period of fastest growth.
 - Supplementation is not necessary, but if desired, other food items and supplements should make up no more than 20% of the total dietary intake.
- Once the adolescent bird is fully feathered, transition to Mazuri® Waterfowl Maintenance Diet. For birds laying eggs, we recommend Mazuri® Waterfowl Layer Diet.
- Always provide plenty of fresh, clean water.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

Storage Conditions

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.