

Mazuri® Sea Duck Diet

(Available at www.mazuri.com or through a Mazuri® retailer)



Formula Code - 5681

Description

Mazuri® Sea Duck Diet is an extruded floating ration formulated for growing, breeding and adult sea ducks and other diving duck species.

Features and Benefits

- **Floating ration** - Floats for a limited period of time, limiting water contamination and waste.
- **Balanced levels of essential nutrients** - Contains high quality protein from animal sources.
- **Sustainable fishmeal source used.**

Product Form

Extruded pellet size: ¼" diameter x ¼" length.

- 50 lb. net weight paper sack

Catalog

0006599



Guaranteed Analysis

Crude protein not less than	21.5%	Calcium not less than	2.25%
Crude fat not less than	5.0%	Calcium not more than	2.75%
Crude fiber not more than	5.0%	Phosphorus not less than	1.20%
Ash not more than	12.0%	Salt not less than	0.25%
		Salt not more than	0.75%
		Sodium not more than	0.55%

Ingredients

Fish meal (menhaden), ground corn, whole wheat, wheat middlings, dehydrated alfalfa meal, porcine meat and bone meal, brewers dried yeast, calcium carbonate, whey, soybean oil, salt, dicalcium phosphate, glyceryl monostearate, magnesium oxide, pyridoxine hydrochloride, dried Lactobacillus acidophilus fermentation product, dried Lactobacillus casei fermentation product, dried Bifidobacterium thermophilum fermentation product, choline chloride, menadione sodium bisulfite complex (source of vitamin K), dried Enterococcus faecium fermentation product, vitamin D₃ supplement, vitamin A supplement, d-alpha tocopheryl acetate (form of vitamin E), biotin, preserved with mixed tocopherols (form of vitamin E), rosemary extract, calcium pantothenate, citric acid, folic acid, thiamine mononitrate, nicotinic acid, riboflavin supplement, vitamin B₁₂ supplement, manganous oxide, zinc oxide, copper sulfate, calcium iodate, sodium selenite.

Feeding Directions

Mazuri® Sea Duck Diet is a floating ration. Feed ad libitum on surface of water. If supplementation is desired, species-appropriate supplementation should be provided at no more than 20% of the total dietary intake (as is basis).