



## High-Fat Hand Feeding Formula

Mazuri® High-Fat Hand Feeding Formula is designed for hand feeding baby birds of fruit- and seed-eating species that would benefit from higher fat levels, such as macaws, African greys, Eclectus, and larger psittacine species.

---

### Features and Benefits

- **Contains probiotics** – For optimized digestion and immune support.
- **Contains omega-3 fatty acids** – Supports healthy development.
- **Higher fat formula** – When compared to Mazuri® Hand Feeding Formula 5ZM3.
- **Formulated to support healthy growth and feather development.**
- **No supplementation required.**
- **No added artificial colors.**

---

### Product Form & Packaging

Meal |

- **Catalog #3009526-745** | 4 lb. (1.81 kg) net weight resealable poly bag
- **Catalog #3009526-746** | Case: 6 x 4 lb. resealable poly bags

---

### Guaranteed Analysis

Crude protein not less than .....	20.00%	Moisture not more than .....	12.00%
Crude fat not less than .....	16.00%	Ash not more than .....	7.00%
Crude fiber not more than .....	2.50%	Omega-3 fatty acids not less than .....	1.00%
Total Microorganisms .....			270,000 CFU/g

(Lactobacillus acidophilus, Lactobacillus casei, Bifidobacterium thermophilum, Enterococcus faecium)\*

\*Contains a source of live (viable) naturally occurring microorganisms.

---

### Ingredients

Wheat Flour, Dehulled Soybean Meal, Soybean Oil, Corn Starch, Soy Protein Concentrate, Sucrose, Dried Egg Product, Brewers Dried Yeast, Dicalcium Phosphate, Calcium Carbonate, Lecithin, Salt, DL-Methionine, Yeast Culture, Choline Chloride, Dried Lactobacillus acidophilus Fermentation Product, Dried Lactobacillus casei Fermentation Product, Dried Bifidobacterium thermophilum Fermentation Product, Spirulina Algae Meal, Dried Enterococcus faecium Fermentation Product, Preserved with Mixed Tocopherols, L-Ascorbyl-2-Polyphosphate (Vitamin C), Citric Acid (a Preservative), Manganese Sulfate, Ferrous Sulfate, Zinc Sulfate, Vitamin E Supplement, Niacin Supplement, Calcium Pantothenate, Canthaxanthin (color), Copper Sulfate, Pyridoxine Hydrochloride, Riboflavin Supplement, Thiamine Mononitrate, Menadione Sodium Bisulfite Complex (Vitamin K), Rosemary Extract, Folic Acid, Potassium Iodate, Vitamin A Supplement, Biotin, Beta Carotene, Sodium Selenite, Vitamin D3 Supplement, Vitamin B12 Supplement.

---

### Feeding Directions

NEVER FEED DRY POWDER

- For best care, establish a planned feeding program using sanitary practices. Mix with water to appropriate consistency for the age of chick that you are feeding. Young chicks will require more water.
  - As chicks grow, feeding frequency will decrease; allow crop to empty before next feeding. Feed until crop is full or chick refuses to eat.
- To ensure complete hydration of the product, mix with hot water and feed when the temperature is between 100° to 105° F (38° to 40.5° C). If it is too hot, add a small amount of cold water to reach feeding temperatures, or wait until the formula reaches an appropriate feeding temperature.



- Water:powder ratio may vary among species. Measuring of powder by weight is recommended. Suggested mixing ratios should be modified to fit the nutritional and hydration needs of the birds you are feeding and to ensure that the product will flow through the syringe you are using.
  - In the first weeks post-hatch, 90% Water: 10% Powder (measure by weight).
  - As the bird grows, 70% Water: 30% Powder (measure by weight).
- Mix thoroughly with a fork or whisk until fully mixed.
  - This product contains pigment ingredients important to the birds. Some pigments may not be fully absorbed, and pink feces may occasionally be observed. This is normal and may vary over time.
- Feed until the crop is full using the appropriate hand-feeding syringe. Do not overfeed.
  - Most chicks will eat 10% or more of their body weight per feeding. Growth rates can vary by bird and species, so monitor body weight daily. This is the most accurate way to ensure proper growth and feeding rates.

#### **Feeding Notes:**

- Under-heating can cause crop stasis, and over-heating can cause crop burn.
- Never use a microwave to heat formula as it can create hot spots and cause crop burns.
- Never feed dry powder. Always mix fresh with water; do not reuse.
- Make a fresh batch of formula for each feeding. Reheating is not advised.
- Do not use formula for more than 6 hours after preparing.
- Do not dilute the formula excessively as this will reduce calories fed to the chick and will impair growth and development.
- Use good sanitation practices and good quality water.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals.
- This diet is not for human consumption.

---

#### **Storage Conditions**

For best results, reseal the bag between uses or store contents of open paper sack in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Use within 1 year of bag manufacturing or "Best if Used By" date.